

(12) UK Patent Application (19) GB (11) 2 394 951 (13) A

(43) Date of A Publication 12.05.2004

(21) Application No: 0225701.2

(22) Date of Filing: 04.11.2002

(71) Applicant(s):
Cipla Limited
(Incorporated in India)
289 Bellasis Road, Mumbai Central,
Mumbai 400 008, India

(72) Inventor(s):
Dharmaraj Ramachandra Rao
Rajendra Narayanrao Kankan

(74) Agent and/or Address for Service:
A A Thornton & Co
235 High Holborn, LONDON, WC1V 7LE,
United Kingdom

(51) INT CL⁷:
C07D 277/82 , C07C 233/32

(52) UK CL (Edition W):
C2C CDL CEJ C1390 C213 C22Y C221 C225 C25Y C256
C281 C30Y C31Y C314 C32Y C321 C338 C34Y C342
C351 C353 C440 C45Y C450 C591 C616 C617 C62X
C630 C670 C69Y C774

(56) Documents Cited:
EP 0186087 A1

(58) Field of Search:
UK CL (Edition V) **C2C**
INT CL⁷ **C07C, C07D**
Other: **ONLINE: EPODOC, WPI, JAPIO, CAS-ONLINE**

(54) Abstract Title: **ONE POT SYNTHESIS OF 2,6-DIAMINO-4,5,6,7-TETRAHYDRO-BENZOTHIAZOLE**

(57) 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which is useful for making pramipexole, is made by: (i) reacting bromine with a solution of 4-acetamido-cyclohexanone in water to produce 2-bromo-4-acetamido-cyclohexanone; (ii) after step (i), adding thiourea to produce 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole; (iii) after step (ii), adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole dihydrobromide; and (iv) after step (iii), isolating 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole.

GB 2 394 951 A

Original Printed on Recycled Paper


Process

This invention relates to a process for making 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, an intermediate useful in the production of pramipexole. The invention also relates to the synthesis of pramipexole.

(S)-4,5,6,7-tetrahydro-N6-propyl-2,6-benzothiazole-diamine (or (S)-2-amino-4,5,6,7-tetrahydro-6-(propylamino)benzothiazole), more commonly known as pramipexole, is used in both early and late Parkinson's disease as a dopamine agonist, to stimulate dopamine receptors in the brain. This has been described in EP 0 186 087.

EP 0 186 087 also describes the synthesis of various tetrahydro benzothiazoles, including pramipexole. In particular, the synthesis of pramipexole by the following reaction pathway is described. An initial reaction between bromine and 4-acetylamido-cyclohexanone is carried out in glacial acetic acid, with stirring for several hours, at room temperature. This is followed by the addition of thiourea under refluxing conditions. The reaction mixture is cooled, and crystals of 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole-hydrobromide are precipitated. The precipitate is filtered, then washed with water and acetone. The crystals are then dissolved in hydrobromic acid and the solution is refluxed for several hours. The solution is then concentrated by evaporation and the residue dissolved in methanol, from which crystals of 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide are formed. Subsequently, the 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide may be converted to pramipexole.

This synthesis is illustrated by the following reaction scheme:



This synthetic pathway involves separate reaction steps, each requiring different conditions, solvents, temperatures, etc. This necessitates a discontinuous process and more than one isolation step, which entails longer processing time, lower yields (product is lost during each isolation step), increased effluent load and increased solvent usage, in comparison with a continuous process.

We have now found a way of synthesising 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole from 4-acetamido-cyclohexanone, which avoids the multiple isolation steps used in the previously described processes.

According to one aspect of the present invention there is provided a method of synthesising 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which method comprises comprising: (i) reacting bromine with a solution of 4-acetamido-cyclohexanone in water to produce 2-bromo-4-acetamido-cyclohexanone; (ii) after step (i), adding thiourea to produce 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole-dihydrobromide; (iii) after step (ii), adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole; and (iv) after step (iii), isolating 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole free base.

It is an important feature of the present invention that step (iii) is carried out without any isolation of the 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole produced in step (ii). Thus the entire synthesis can be carried out in a single reaction vessel. Preferably at least three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.

Prior to step (i), the method may comprise the step of oxidising 4-acetamido-cyclohexanol to produce 4-acetamido-cyclohexanone. This step may be carried out in the same reaction vessel as subsequent steps (i) to (iv), thereby avoiding an additional isolation step.

The oxidation reaction may be carried out using oxidising agents including, for example, Jones reagent, sodium hypochlorite, manganese dioxide, pyridinium dichromate or potassium permanganate.

In step (i), the 4-acetamido-cyclohexanone solution and the bromine are preferably combined in the reaction vessel at a temperature in the range 5°C to 75°C, more preferably in the range 15°C to 40°C, and most preferably about room temperature (approximately 25°C). The bromine is preferably added dropwise to the 4-acetamido-cyclohexanone solution. After the bromine and the 4-acetamido-cyclohexanone solution have been combined, the mixture is preferably heated to a temperature in the range 30°C to 80°C, more preferably 40°C to 50°C,

and most preferably about 45°C, and maintained at or near this temperature until the bromination is complete. The completion of bromination is indicated by the elimination of the characteristic brown colour of the bromine.

In step (ii), the temperature is preferably increased to 50°C to 95°C, more preferably to 70°C to 90°C, and most preferably to about 80°C.

In step (iii), the reaction mixture is preferably refluxed.

In step (iv), the reaction mixture is preferably cooled to 1°C to 35°C, more preferably to 5°C to 20°C, and most preferably to about 10°C, and the mixture is then neutralised. Typically the neutralisation is carried out with caustic lye solution (NaOH), although other alkalis may be used. Following neutralisation, the product, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, is isolated. The isolation may be carried out by filtration, centrifugation or any other suitable means. Following isolation, the product is preferably washed with chilled water.

The starting compound, 4-acetamido-cyclohexanone, may conveniently be formed by the oxidation of 4-acetamido-cyclohexanol.

The above described compound, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, is useful as an intermediate in the production of pramipexole and related compounds.

According to another aspect of the present invention there is provided a method of synthesising pramipexole, comprising the steps of : forming 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole by the method of the present invention, then converting it to pramipexole.

The conversion of 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole to pramipexole is well known in the prior art and is described, for example, in US 4,731,374. Any of the methods described in US 4,731,374, may be used in the present invention.

In one embodiment, the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole is converted to pramipexole by reaction with a propionyl halide, such as propionyl bromide, under suitable reaction conditions.

Both 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole and pramipexole have an asymmetric carbon atom, and exist as two distinct enantiomers: the S(-) isomer and the R(+)

isomer. The pharmacological activity of the S(-) isomer of pramipexole is, however, twice as high as that of the R(+) isomer, and the name pramipexole is commonly used to refer to the optically pure S(-) form. In this specification, "2,6-diamino-4,5,6,7-tetrahydro-benzthiazole" encompasses the R(+) and S(-) enantiomers individually and also encompasses any mixture thereof including the racemic mixture, and the term "pramipexole" encompasses the R(+) and S(-) enantiomers individually and also the racemic mixture.

The resolution of a racemic mixture of 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole can be carried out after step (iv) above. Methods of resolution are known in the art. Alternatively, pramipexole racemate can be produced prior to resolution, then the mixture resolved, if desired.

The resolution of pramipexole racemate is described by Schneider and Mierau (J. Med. Chem. 30, 494 (1987)). This method uses the di-amino derivative of (\pm)-4,5,6,7-tetrahydro-N6-propyl-2,6-benzothiazolodiamine as a substrate and L(+) tartaric acid as a resolution agent. Following resolution, optically active pramipexole has been prepared by two-step propylation of the single enantiomer of the di-amino precursor comprising the steps of reaction with propionanhydride followed by a reduction of the propionyl intermediate.

The synthesis of the present invention avoids the need to isolate intermediate compounds, and thus the yield is higher and the processing time is lower. Furthermore, owing to the absence of an organic solvent (such as acetic acid) the costs are lower, and the reaction conditions are milder – the milder reaction conditions also have a positive impact on product purity.

The invention will now be further described with reference to the following Examples.

Example 1

Bromine (112g) was added dropwise to a solution of 4-acetamidocyclohexanone (100g) in 500ml water at room temperature. The mixture was warmed to approximately 45°C and maintained at this temperature until the bromine colour had been lost. To this,

thiourea (125g) was added, and the mixture was heated to approximately 80°C. To this, aqueous hydrobromic acid (100ml) was added, and the contents of the reaction vessel were refluxed. The contents were then cooled to approximately 10°C, and neutralized with caustic lye solution. The product, 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole, was isolated by filtration, and washed with chilled water and dried. The product was off-white in colour, and the yield was approximately 60g in weight.

Example 2

To a solution of 4-acetamido-cyclohexanol (100 g) in acetone (1L) was added Jones reagent (prepared from 68.5 g chromic oxide, 105 g sulphuric acid and 400 ml water) at 10-15°C. The excess of the reagent was quenched by addition of isopropanol (400 ml) and the solvent was removed under reduced pressure. Ethyl acetate (600 ml) was added, the contents stirred for 10 minutes and the lower aqueous portion drained off. Ethyl acetate was concentrated under reduced pressure and the residue dissolved in water (500 ml). Bromine (112 g) was added dropwise and the further reactions were carried out as described in Example 1.

Example 3

To a suspension of 4-acetamido-cyclohexanol (100 g) in water (300 ml) was added a solution of 10% sodium hypochlorite (500ml) and the contents stirred at room temperature for 12 hours. To this was added liquid bromine (112 g) and further reactions carried out as described in Example 1.

It will be appreciated that the invention described above may be modified.

CLAIMS:

- 1 A method of making 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole, which method comprises the steps in sequence of: (i) reacting bromine with a solution of 4-acetamidocyclohexanone in water to produce 2-bromo-4-acetamidocyclohexanone; (ii) adding thiourea to produce 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole; (iii) adding an aqueous solution of hydrobromic acid to produce 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole; and (iv) isolating 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole.
- 2 A method according to claim 1 wherein step (iii) is carried out without isolating the 6-acetylamino-2-amino-4,5,6,7-tetrahydro-benzthiazole produced in step (ii).
- 3 A method according to claim 1 or 2, wherein any three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.
- 4 A method according to claim 1, 2 or 3, wherein steps (i) to (iv) are carried out in a single reaction vessel.
- 5 A method according to claim 1, 2, 3 or 4, further comprising, prior to step (i), the step of oxidising 4-acetamidocyclohexanol to produce 4-acetamidocyclohexanone.
- 6 A method according to claim 5, wherein the step of oxidising 4-acetamidocyclohexanol to produce 4-acetamidocyclohexanone and at least three successive steps of steps (i) to (iv) are carried out in a single reaction vessel.
- 7 A method according to any preceding claim wherein in step (i) the solution of 4-acetamidocyclohexanone in water and bromine are combined at a temperature of from 15°C to 40°C.

- 8 A method according to any preceding claim wherein, after the bromine and the 4-acetamido-cyclohexanone solution have been combined, the mixture is heated to a temperature of from 40°C to 50°C, and maintained at or near this temperature until the bromination is complete.
- 9 A method according to any preceding claim wherein, in step (ii), the temperature is increased to 70°C to 90°C.
- 10 A method according to any preceding claim, wherein step (iii) is carried out under refluxing conditions.
- 11 A method according to any preceding claim wherein, after step (iii) but before step (iv), the reaction mixture is cooled to 5°C to 20°C, then neutralised.
- 12 A method according to any preceding claim, further comprising the step of resolving the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole isolated in step (iv) into its R(+) and S(-) enantiomers and recovering the R(+) and/or S(-) enantiomer.
- 13 A method of synthesising pramipexole, comprising the steps of : forming 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole by a method according to any preceding claim, and converting it to pramipexole.
- 14 A method according to claim 13, wherein 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole is converted to pramipexole by reaction with a propionyl halide.
- 15 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises the R(+) enantiomer.

- 16 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises the S(-) enantiomer.
- 17 A method according to claim 13 or 14, wherein the 2,6-diamino-4,5,6,7-tetrahydro-benzothiazole comprises a racemic mixture.
- 18 A method according to claim 14, further comprising the step of resolving the pramipexole into its R(+) and S(-) enantiomers and recovering the R(+) and/or S(-) enantiomer.
- 19 A method of synthesizing 2,6-diamino-4,5,6,7-tetrahydro-benzthiazole substantially as herein described with reference to the Examples.



INVESTOR IN PEOPLE

Application No: GB 0225701.2
Claims searched: 1-19

Examiner: Dr. Simon Grand
Date of search: 1 April 2003

Patents Act 1977 : Search Report under Section 17

Documents considered to be relevant:

Category	Relevant to claims	Identity of document and passage or figure of particular relevance
A		EP 0186087 A1 (THOMAE) See example 3

Categories:

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

Field of Search:

Search of GB, EP, WO & US patent documents classified in the following areas of the UKC^v:

C2C

Worldwide search of patent documents classified in the following areas of the IPC^v:

C07D and C07C

The following online and other databases have been used in the preparation of this search report:

ONLINE: EPODOC, WPI, JAPIO, CAS-ONLINE